

Leinen Health Products

Exhibits

012 3039

Drug Facts

Active Ingredients (in each caplet):
 Acetaminophen 500 mg
 Diphenhydramine Hydrochloride 25 mg
Purposes: Pain Reliever/Fever Reducer Sleep Aid

Uses • For the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness.

Warnings
 Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.
 Do not use • with other products containing acetaminophen
 • if you have a breathing problem such as emphysema or chronic bronchitis
 • if you have glaucoma or difficulty in urination due to enlargement of the prostate gland.
 Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product • do not exceed recommended dosage
 • drowsiness will occur
 • Do not drive a motor vehicle or operate machinery
 • Do not use for more than 10 days • Avoid alcoholic beverages

NDC 59606-060-24

equate

extra strength pm

NIGHTTIME SLEEP AID/PAIN RELIEVER

150 CAPLETS Compare to Extra Strength Tylenol® PM
 Caplets active ingredients*

EQ81129AQ08
LHP016

LHP016

NDC 59606-060-24

equate

extra strength pm

NIGHTTIME SLEEP AID/PAIN RELIEVER

150 CAPLETS Compare to Extra Strength Tylenol® PM
 Caplets active ingredients*

USE ONLY IF INNER FOIL SEAL PRINTED "SEALED FOR YOUR PROTECTION" IS NOT BROKEN OR MISSING. Keep tightly closed. Store at room temperature between 15° and 30° C (59° and 86° F). Use by expiration date printed on package.

* This product is not manufacturer distributed by McNeil Consumer Products Company, distributor of Extra Strength Tylenol, PM. Tylenol is a registered trademark of Ortho-Clinical Diagnostics, Inc.

MANUFACTURED BY
LEINER HEALTH PRODUCTS INC.
CARSON, CA 90745 U.S.A.
EQ81129AQ08

Made in the USA
SATISFACTION GUARANTEED
BY REFUND OR EXCHANGE

Drug Facts (continued)

Stop use and ask a doctor • if this product is used for pain for more than 10 days
 • if fever persists for more than 3 days or gets worse
 • if redness or swelling is present, these could be signs of a serious condition
 • if symptoms persist or new ones occur
 • if sleeplessness persists continuously for more than 2 weeks, insomnia may be a symptom of serious underlying medical illness.

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions Adults and children 12 years of age and over: 2 caplets at bedtime. Do not give to children under 12 years of age.

Inactive Ingredients: FD&C Blue No. 1, FD&C Blue No. 2, Hydroxypropyl Methylcellulose, Microcrystalline Cellulose, Polyethylene Glycol, Triphenyl Methane Dioxide, and may also contain: Croscarmellose Sodium, Magnesium Stearate, Maleic Acid, Polydextrose, Polyethylene Glycol, Polyethylene Glycol, Povidone, Pregelatinized Starch, Silicon Dioxide, Sodium Starch Glycolate, Stearic Acid, Tracetin.



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Federal medical attention is critical for adults as well as children. Do not use if you do not notice any signs or symptoms.

Directions

| Age Group | Frequency | Maximum Dose |
|---|-------------------------------------|--------------------------------------|
| Adults and children 12 years of age and over: | 2 softgels with water every 4 hours | Not to exceed 8 softgels in 24 hours |
| Children 8 to under 12 years of age: | 1 softgel with water every 4 hours | Not to exceed 4 softgels in 24 hours |
| Children under 8 years of age: | Consult a doctor | |

Inactive Ingredients: D&C Yellow No. 10, Edible Ink, FD&C Red No. 40, FD&C Yellow No. 6, Gelatin, Glycerin, Polyethylene Glycol, Povidone, Propylene Glycol, Propyl Paraben, Purified Water, Sorbitol

Store at room temperature between 15° and 30°C (59° and 86°F). Use by expiration date printed on package. Keep tablets in carton for better identification until used.

*This product is not manufactured or distributed by Procter & Gamble, distributor of DayQuil® LiquidCaps® DayQuil® and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

CF81744A008
 Distributed by:
 Lannet Health Products Inc.
 Carson, CA 90245 1-800-774-2676
 Made in the U.S.A.



IFC236

96 SOFTGELS

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

- Cough Suppressant
- Pain Reliever - Fever Reducer
- Nasal Decongestant

Pseudoephedrine / Acetaminophen / Dextromethorphan

Cold/Flu Medicine Softgels

DAY-TIME
Non-Drowsy

ITEM # 314005

KIRKLAND

96 SOFTGELS

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

- Cough Suppressant
- Pain Reliever - Fever Reducer
- Nasal Decongestant

Pseudoephedrine / Acetaminophen / Dextromethorphan

Cold/Flu Medicine Softgels

DAY-TIME
Non-Drowsy

KIRKLAND

96 SOFTGELS

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

- Cough Suppressant
- Pain Reliever - Fever Reducer
- Nasal Decongestant

Pseudoephedrine / Acetaminophen / Dextromethorphan

Cold/Flu Medicine Softgels

DAY-TIME
Non-Drowsy

KIRKLAND

96 SOFTGELS

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

Compare To The Active Ingredients Of DayQuil® LiquidCaps™

- Cough Suppressant
- Pain Reliever - Fever Reducer
- Nasal Decongestant

Pseudoephedrine / Acetaminophen / Dextromethorphan

Cold/Flu Medicine Softgels

DAY-TIME
Non-Drowsy

KIRKLAND

KIRKLAND

Non-Drowsy
DAY-TIME
 Cold/Flu Medicine Softgels
 Pseudoephedrine / Acetaminophen / Dextromethorphan

- Nasal Decongestant
- Pain Reliever - Fever Reducer
- Cough Suppressant

SAFETY SEAL: THIS PRODUCT IS PROTECTED WITH SEALED BUSTER UNITS. DO NOT USE IF ANY ARE DAMAGED OR MISSING.

Drug Facts

| Active Ingredients (in each softgel) | Purpose |
|--------------------------------------|-----------------------------|
| Pseudoephedrine HCl 30 mg | Nasal Decongestant |
| Acetaminophen 250 mg | Pain Reliever/Fever Reducer |
| Dextromethorphan HBr 10 mg | Cough Suppressant |

Uses temporarily relieves symptoms due to the common cold/flu:
 • fever, sore throat, pain, • headache • muscular aches
 • sore throat, pain • nasal congestion • cough

Warnings
 Alcohol Warning: If you consume 3 or more alcoholic drinks each day, do not use this product until you are no longer intoxicated. Alcohol consumption may cause low blood pressure.
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's disease, or high blood pressure) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a health professional before taking this product. • with other products containing acetaminophen.

Ask a doctor before use if you have:
 • asthma • progressive phlegm (mucus)
 • chronic bronchitis • heart disease • persistent or chronic cough • emphysema
 • cough associated with smoking • high blood pressure • thyroid disease
 • diabetes • difficulty in urination due to enlargement of the prostate gland

See use and ask a doctor if:
 • if nervousness, dizziness, or sleeplessness occur
 • if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting
 • if symptoms do not improve or are accompanied by fever that lasts more than 3 days, or if new symptoms occur.
 • if cough persists for more than 7 days (adults) or 5 days (children), recurs, or is accompanied by rash or persistent headache. A persistent cough may be a sign of a serious condition.

Failure to follow these warnings could result in serious consequences.
 Respirator (Breathers): Do not use for more than 7 days (for adults) or 5 days (for children).


Exhibit B

Drug Facts

Purposes
Acetaminophen 500 mg
Pain Reliever/Fever Reducer

Active Ingredients (in each caplet)
Acetaminophen 500 mg

Warnings
USPS • temporarily relieves minor aches and pains associated with: cold, flu, headache, toothache, muscular aches, backache, the premenstrual and menstrual periods (dysmenorrhea), fever, arthritis, rheumatism • and to reduce fever
Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers.
Acetaminophen may cause liver damage.
Do not use • with other products containing acetaminophen.



Member's Mark™

Pain Reliever • Fever Reducer

ACETAMINOPHEN

500 mg

Compare to Extra Strength Tylenol® active ingredient

Non Aspirin

EXTRA STRENGTH

500 Caplets

100

Drug Facts (continued)

Stop use and ask a doctor if this product is used for pain for more than 10 days or for swelling is present, these could be signs of a serious condition.
• if pain or fever persists or gets worse • if new symptoms occur • or if redness or if pregnant or breast-feeding, ask a health professional before use.
Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.


Directions

| Age Group | Frequency | Maximum Dosage |
|---|-------------------------|-------------------------------------|
| Adults and children 12 years of age and over: | 2 caplets every 6 hours | Not to exceed 8 caplets in 24 hours |
| Children under 12 years of age: | Consult a doctor | |

Inactive Ingredients: Hydroxypropyl Methylcellulose, Povidone, Corn Starch, and Stearic Acid. May also contain: Carnauba Wax, Croscarmellose Sodium, Hydroxypropyl Cellulose, Polyethylene Glycol, Sodium Starch Glycolate, Talc, and Titanium Dioxide.

USE ONLY IF INNER FOIL SEAL PRINTED - SEALED FOR YOUR PROTECTION IS NOT BROKEN OR MISSING.
Keep tightly closed. Store at room temperature between 15° and 30°C (59° and 86°F). Use by expiration date printed on package.
Distributed by: SWC
608 S.W. 8th Street, Bentonville, AR 72712
MM98080M06

Lot No. Exp. Date.



Leiner Health Products 810 E. 233 STREET • CARSON, CA 90745
In-house Graphics Department VOICE (310) 835-8400 • FAX (310) 522-9180

Software(s): Illustrator 8 MAC NO
 IBM MAC

Die Number: 3 X 8
Prepared by: nance
 01 17 01

Job No. 0488-1000248 **Proofer:**

COLORS

| | | | | | | | |
|---------------|--------------|----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Process Black | Process Cyan | Process Yellow | Process PMS 186 | Process PMS 872 | Process PMS 000 | Process PMS 000 | Process PMS 000 |
|---------------|--------------|----------------|-----------------|-----------------|-----------------|-----------------|-----------------|

Free Penet: Universe, Lucida Sans, ITC Stone Serif

Ingredient Panels: Helvetica Black, Helvetica Condensed, Helvetica

Smallest Point Size: 5 pt

Path: PRODUCTION\NOTCAMEMBERS\MARK\ARTWORK\LABELS\MM98080B.EPS

CAUTION:
It is the vendors responsibility to print a scannable UPC code from either our artwork or your alternative UPC method.

*COLOR BREAK AND DIE LINES ARE FOR REFERENCE ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRINT PRODUCTION.
PMS 186 REPLACES MAGENTA

USE ONLY IF INNER FOL SEAL PRINTED "SEALED FOR YOUR PROTECTION" IS NOT BROKEN OR MISSING.

INDICATIONS: For temporary relief of the pain of headache, sinusitis, colds, muscular aches, menstrual discomfort, toothaches and minor arthritis pain.

DIRECTIONS: Adults and children 12 years of age and over: 2 tablets every 6 hours while symptoms persist, not to exceed 8 tablets in 24 hours or as directed by a doctor. Drink a full glass of water with each dose. Children under 12 years of age: Consult a doctor.

Warnings: Children and teenagers who have or are recovering from chicken pox, the symptoms, or flu should NOT use this product. If nausea, vomiting, or fever occur, consult a doctor because these symptoms could be an early sign of Reye syndrome, a rare but serious illness. Do not take this product for pain for more than 10 days or for fever for more than 3 days. If pain persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor.

WARNINGS: Children and teenagers who have or are recovering from chicken pox, the symptoms, or flu should NOT use this product. If nausea, vomiting, or fever occur, consult a doctor because these symptoms could be an early sign of Reye syndrome, a rare but serious illness. Do not take this product for pain for more than 10 days or for fever for more than 3 days. If pain persists or gets worse, if new symptoms occur, or if redness or swelling is present, consult a doctor.

Do not take this product if you are allergic to aspirin or if you have asthma unless directed by a doctor. If ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product. Do not take this product if you have stomach problems (such as heartburn, upset stomach, or stomach pain) that persist or recur, or if you have ulcers or bleeding problems, unless directed by a doctor. Do not take this product if you are pregnant or nursing a baby, with any drug, if you are pregnant or nursing a baby, or if you have any other medical condition. Ask your doctor or pharmacist for more information. Do not use this product if it is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

SAFEWAY

PAIN RELIEVER



NDC 21130-310-14

COMPARE TO ACTIVE INGREDIENTS IN EXCEDRIN[®]

Extra Strength Pain Relief Tablets

• acetaminophen, aspirin and caffeine

50 COATED TABLETS

SAFEWAY

PAIN RELIEVER



NDC 21130-310-14

COMPARE TO ACTIVE INGREDIENTS IN EXCEDRIN[®]

Extra Strength Pain Relief Tablets

• acetaminophen, aspirin and caffeine

50 COATED TABLETS

SAFEWAY

PAIN RELIEVER



NDC 21130-310-14

COMPARE TO ACTIVE INGREDIENTS IN EXCEDRIN[®]

Extra Strength Pain Relief Tablets

• acetaminophen, aspirin and caffeine

50 COATED TABLETS

ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen and aspirin or other pain relievers/fever reducers. Acetaminophen and aspirin may cause liver damage and stomach bleeding.

DRUG INTERACTION PRECAUTION: Do not take this product if you are taking a prescription drug for anti-coagulation (thinning the blood), diabetes, gout, or arthritis unless directed by a doctor.

ACTIVE INGREDIENTS: Each tablet contains: Acetaminophen 250 mg., Aspirin 250 mg., Caffeine 65 mg.

INACTIVE INGREDIENTS: Hydroxypropyl Methylcellulose, Polyethylene Glycol, Stearic Acid. May also contain: Carnauba Wax, Cellulose, FD&C Blue No. 1, Hydroxypropyl Cellulose, Methylcellulose, Macrogol 400, Polyethylene Glycol, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Starch, Triammonium Citrate, and Cellulose.

Keep tightly closed. Store at room temperature between 15 and 30°C (59 and 86°F). Use by expiration date printed on package.

We appreciate your comments and ideas. Write to us or call toll free 1-800-SAFEWAY. Please include product code and lot number. www.safeway.com

QUARTER SERRATION

PRODUCT OF U.S.A.

DISTRIBUTED BY SAFEWAY INC. CA 94568-0009

SW95801BA08

Squibb Company.

Bristol-Myers Products, distributor of Excedrin[®]. Excedrin[®] is a registered trademark of Bristol-Myers.

This product is not manufactured or distributed by Bristol-Myers.

25% BLACK OVERPRINTS
SOLID PMS 485

LHP010


SW95801BA08
LHP010

SAFEWAY

Extra Strength Pain Relief Tablets

50 COATED TABLETS

TAMPER EVIDENT BOTTLE
DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.



Leiner Health Products
In-house Graphics Department

810 E. 233 STREET • CARSON, CA 90745
VOICE (310) 835-8400 • FAX (310) 522-9180

Software(s):
Illustrator 8.0

WamNet:
 YES
 NO

IBM MAC NO

COLORS

Pantone Black, Pantone Pro Yellow, PMS 485, PMS 343, PMS 000, PMS 000, PMS 000

Fonts

Face Panel: Times, Helvetica Condensed

Ingredient Panels: Helvetica, Helvetica Black

Smallest Type: 5.5 pt

Die Number: LHP010

Prepared by: Lois 11-30-00

Path File Name: Q:\PRODUCTION\MTGSAFEWAY\ARTW\

SW-1000334 Proofreader:

*COLOR BREAK AND DIE LINES ARE FOR REF. Exhibit E ATLELY REFLECT ACTUAL PRINT PRODUCTION.

the manufacturer's responsibility for a complete UPC code in all our artwork or alternative UPC method.